

LODE GOLD OCCURRENCES NEAR
THE KAKO AND STUYAHOK
PLACER MINES, SOUTHWESTERN
ALASKA

GEOLOGIC REPORT 1/10/90

Bruce Hickok, June McAtee
Geologists
CALISTA CORPORATION
601 West Fifth Avenue
Anchorage, Alaska 99501
(907) 279-5516

TABLE OF CONTENTS

KAKO AND STUYAHOK GEOLOGIC REPORT

SUMMARY.....	3
LOCATION AND ACCESS.....	4
LAND.....	4
REGIONAL GEOLOGY.....	5
TECTONIC AND STRUCTURAL SETTING.....	6
REGIONAL MINERALIZATION.....	6
KAKO HISTORY.....	7
KAKO GEOLOGY.....	8
KAKO PLACER	10
LODE TARGETS.....	11
STUYAHOK HISTORY.....	14
STUYAHOK GEOLOGY.....	14
STUYAHOK PLACER.....	15
LODE TARGETS.....	15
CONCLUSIONS.....	16
RECOMMENDATIONS.....	16

Summary

The Kako and Stuyahok placer mines in southwestern Alaska have a geologic setting conducive to the discovery of a deep level epithermal rhyolite hosted or associated gold system. The Kako and Stuyahok occurrences are related to rhyolite dikes and stocks which intrude the sediments and volcanics topographically above and directly beneath the placer deposits. While no mineable areas of ore grade mineralization have yet been discovered, these placer mines both have large areas of bedrock which are anomalous in gold, silver, arsenic, and mercury. There are indications of structural control on the location and geometry of the occurrences found to date.

The gangue minerals, angularity and shape of the gold, and the volume of cinnabar, arsenopyrite and stibnite in the placer concentrates are all consistent with placer deposits derived from an epithermal lode environment. Significant placer gold resources remain in the Kako and Stuyahok camps.

The areas are within 6 and 20 miles of the Yukon river respectively. Stuyahok is roughly 15 miles northeast of Kako. Good winter haul roads are already in existence. Access to this area rivals most areas of Alaska.

KAKO

At Kako large areas of the ridge above the placer deposits have been shown to be anomalous in gold and arsenic. A 1000 feet by 2000 feet soil and rock chip grid produced an anomaly which is open on two sides with values ranging from 200 to 500 ppb gold and 100 to 2000 ppm arsenic. The anomalous values are related to the contacts of rhyolite stocks and dikes which intrude a chert unit. A rhyolite breccia zone 40 feet wide and at least 1000 feet long traversing the grid has values up to 900 ppb gold. In addition to the anomalous areas on the ridges above the placer camps, anomalous gold, silver, arsenic and mercury values are associated with poorly exposed rhyolite dikes and sills in the valleys, near and under the placer deposits.

STUYAHOK

At Stuyahok rock and stream sediment samples have defined a large area anomalous in gold, silver, arsenic and mercury. Areas of quartz eye rhyolite porphyry and altered Gemuk group volcanics have rock values ranging up to 1 ppm gold and stream sediment samples ranging up to 10 ppm gold.

It is likely that additional mineralized areas will be discovered in

the Kako and Stuyahok placer camps. As a result of very basic work, a lode occurrence has been found within nearly every creek or gulch which hosts a placer deposit. Reconnaissance level sampling has defined large areas of anomalous gold, mercury, antimony and arsenic which suggest large gold systems. The strong arsenic halos which surround gold mineralization at both Kako and Stuyahok are readily detectable. The active placer operations in the two camps will provide support for lode exploration and hopefully in the course of operations will uncover occurrences buried in the valleys.

Location and Access

Location

The Kako and Stuyahok areas are located in western Alaska just north of Yukon River, see Figure 1. The prospects are within 6 and 20 miles of the river respectively and are located close to the villages of Marshall, Russian Mission, and Holy Cross. These placer mines form the heart of the Marshall Mining District.


Access

Access to the Kako property is gained by ocean-going barge up the lower Yukon River and by 4 miles of winter haul road from the landing just above the town of Russian Mission. The road has recently been upgraded to handle summer use. Air access is facilitated by any of several well maintained and not so well maintained airports in the area, including a Hercules size strip (greater than 4,500 feet) at the town of Marshall and 3000 foot long airstrips at Russian Mission and Holy Cross which are serviced by at least twice daily commuter flights. A well maintained 2000 foot strip is located on the Kako property.

Access to the Stuyahok Property is similar to that of Kako, although the winter road is roughly 30 miles long. The airstrip at the placer mine is a casually maintained 1500 feet in length. Both Kako and Stuyahok are served from the regional aviation centers located at St Marys (60 miles from Kako), Aniak (80 miles from Kako), and Bethel (80 miles from Kako). All three are served at least twice daily by jet service from Anchorage.

Land

Calista Corporation is one of twelve regional native corporations established by the Alaska Land Claims Settlement Act. Calista will, when the land selection process is complete, have over seven million acres of subsurface estate. Currently Calista Corporation owns 300 square miles of land holdings in the lower Yukon mineral belt, including many prospects and access corridors.



Calista's land selections were based on village needs and on extensive exploration conducted by Resource Associates of Alaska (RAA) during 1973-1975. Calista still has land selection rights to additional lands and much of the surrounding federal lands are closed to mineral entry pending additional or amended selections. The net result is that the major portion of the mineral potential in Southwestern Alaska is either directly or indirectly controlled by Calista Corporation.

Regional Geology

Terranes

The Lower Yukon area is comprised of two tectonic terranes which were described by Cady, 1955, and Hoare and Coonrad, 1959, these include:

1) A Paleozoic to middle Mesozoic continental margin rift and associated geosynclinal deposits (the Nyac and Gemuk Groups). The Gemuk Group as defined by Hoare and Coonrad, 1959 is comprised of mafic lava flows, breccia, tuffs, and agglomerate, minor felsic flows and tuffs interbedded with siliceous siltstone, chert, graywacke. The Lower Yukon Greenstone Belt is an informal name given to this sequence which covers about 1000 square miles of the Lower Yukon. Most of the Kako and Stuyahok drainage is considered to be Gemuk Group.

And 2) a marine successor basin (the Kuskokwim Group), which followed a major Cretaceous orogeny, that was filled with flysch by the early Tertiary. The Kuskokwim Group in the Lower Yukon is composed mostly of graywacke with minor shale. Only a small part of the Stuyahok drainage is thought to be Kuskokwim group, see Figure 2 and attached map - Geologic and Geochemical Map of the Marshall Area.

Igneous Rocks

Large volumes of upper Tertiary bimodal magma intrude both tectonic terrains. The emplacement was structurally controlled by large displacement right lateral fault systems. The principle igneous rocks are monzonitic stocks, hypabyssal and extrusive albite rhyolites, and biotite basalts. Late Pliocene flood basalts are found south and west of Kako and Stuyahok.

Whole rock analyses from Cady, 1955, Bundtzen, 1987, and unpublished Calista data are consistent with a classification of co-magmatic alkali-calcic, metaluminous volcano-plutonic complexes and related but typically distinct peraluminous rhyolite sills. The volcano-plutonic complexes in southwestern Alaska are part of an extensive belt of reduced (ferrous/ferric ratio) magma as defined by







FIGURE 2

GEOLOGICAL MAP OF LOWER YUKON GOLD BELT

EXPLANATION



CROSS-SECTION LOCATION



Keith, 1988.

The monzonitic stocks are quite variable in composition and frequently have silica-poor intrusive phases interrelated with quartz-rich phases. Whole rock analyses exhibit a variability from alkaline to calc-alkaline monzonitic stocks with an average sub alkaline composition. The derivation and intrusive sequence may be crudely similar to that proposed for Lower Cretaceous and Early Tertiary igneous rocks of the western United States (Bundtzen, 1987).

Tectonic and Structural Setting

Major Faults

In southwestern Alaska the large displacement right lateral fault systems like the Iditarod, Aniak, Holokuk, Holitna, and Farewell, localize Tertiary plutonism. The many fault splays and extensional zones (simoids) between splays and the major faults also controlled the emplacement of igneous rocks. The Kako and Stuyahok areas are influenced by the large displacement Chirosky Fault to the west and the Aniak-Thompson Fault to the east (Hoare and Conrad, 1959), see Figure 3. In the area between the fault systems (Kako and Stuyahok) an extensional tectonic environment was created during the Tertiary and Early Quaternary. The rhyolites of interest were emplaced into this setting.

The area also straddles the Nunivak Arch or Geanticline, a Mesozoic? feature variously interpreted as an orogenic welt or a collision related suture (Box, 1985; Patton, 1973), see Figure 3. In any case, the Paleozoic and Mesozoic rocks adjacent to the arch are characterized by imbricate thrust sheets, isoclinal folds, and sharp, local metamorphic gradients related to deformation or thrust sheet juxtaposition.

Regional Mineralization





Placer Occurrences



Coarse gold placers occur where Tertiary rhyolite or Tertiary-Cretaceous intermediate composition intrusives cut across the sequence described above, see Figure 2. Fine gold placer deposits are reported in many drainages and are not dependent on the occurrence of Cretaceous or Tertiary intrusives.

Lode Occurrences

Lode gold, mercury, and antimony occurrences are spatially associated with the Tertiary igneous rocks. Throughout southwestern Alaska, the alkali olivine quartz monzonite stocks and the peraluminous rhyolites host a variety of gold occurrences which

EXPLANATION

-  SUBAERIAL VOLCANICS
-  RHYOLITE DIKE SWARM
-  QTZ. MONZONITE
-  FAULT



TERTIARY GEOLOGIC MAP
CALISTA REGION

Figure 3

seem to be controlled by emplacement depth, erosional level, and lateral zonation.

The occurrences examined to date generally fall into the following categories: 1. gold-arsenopyrite-scheelite veins and stockworks in monzonites; 2. tourmaline-silver-tin-minor gold greisens in monzonites; 3. low value vein and disseminated gold halos in the hornfels surrounding the stock or dike; 4. gold-stibnite-arsenopyrite-cinnabar veins and micro-stockworks in monzonites, rhyolites and graywacke; and 5. disseminated gold-arsenopyrite-stibnite-cinnabar mineralization in graywacke and rhyolite.

Kako History

Placer gold was discovered at Buster and Bobtail Creek in 1920 by prospectors from the Willow Creek/Marshall area (Cobb, 1973). The gold bearing gravels were mined by hand methods including shaft and adit construction in frozen ground. In the Thirties the Yukon Mining Company used gin poles and scrapers and later a dragline to mine Buster and Bobtail Creeks, Figure 4. By World War 2 the operation was in decline. A rough estimate of 25,000 ounces of gold were produced prior to the 1940's.

In 1974 and 1975 geologists, working for Resource Associates of Alaska (RAA) which was under contract to Calista Corporation to evaluate the mineral potential of the Calista Region, camped at Kako and documented a gold, silver and arsenic anomaly associated with a rhyolite breccia. A large scale rock and soil grid was sampled late in the season of 1974. The results indicated an extensive area of anomalous rhyolite.

RAA's interpretation of the data was that Kako was very promising for both "Nevada style hydrothermal-fracture systems" and "volcanogenic massive sulfide or exhalite" deposits. Many of the soil and stream sediments samples taken by RAA focused on base metal mineralization, often gold and low temperature metals were not tested.

Four Federal placer claims on Buster Creek were kept active and were exempted from the selections made by Calista and the village of Russian Mission following the passage of the Alaska Native Claims Settlement Act. In the early 1980's Dave Penz acquired those four claims on Buster Creek and has built a modern family placer mine featuring a sluice and jigs, a D-8, a one yard hoe, a loader, and other miscellaneous equipment. Penz has produced somewhere around one thousand ounces. Penz received patent on the four claims this in July, 1989.




Fig.4 GEOLOGY AND PLACER CREEKS OF KAKO PLACER DISTRICT

Calista conducted reconnaissance mapping and sampling of the area during a few weeks of the 1985, 1986, 1987 and 1988 field seasons. In 1989 Calista collected over 400 soil and rock chip samples from grids covering areas surrounding the placer deposits. The soil sampling increased the area of anomalous rhyolite and identified several bodies of rhyolite which extend or occur in the placer valleys.

In addition Calista re-assayed 265 selected pulps from RAA's 1974 soil grid program. The pulps were re-analyzed by Bondar-Clegg which used reliable assay methods with lower detection limits than that used by RAA's in house laboratory in 1974. These results defined several new anomalies.

Kako Geology

Lithologic Units


The rocks within the placer drainage are, from south to north, quartz monzonite, rhyolite, chert, and siltstone. Each is in contact with an enclosing andesite flow, agglomerate and tuff complex, Figure 4, Figure 5 and map labeled Geology of Kako Prospect.

Sediments


The Gemuk group sediments exposed on ridges above the Kako placer mine on Buster Creek include cherts, recrystallized limestone, silicified limestone, siliceous siltstones, siliceous shales, and jasper. Directly above the placer creeks the principal sediment is chert.

Cherts in the grid area vary widely. The outcrops at the top of the ridge are white to pale gray, siliceous, and appear amorphous. Most of the cherts seen in rubblecrop are dark gray to greenish color and have a tuffaceous character. At the saddle at the northern end of the grid, the cherts become sedimentary in character, dominated by cherty siltstone with minor cherty agglomerates and gritstones. Some black chert fragments and gray chert with black healed fractures occur as float along lines 500North to 600North. The chert is often fractured and pervasively iron-stained.

The "chert" is variable in appearance and has been mapped in a variety of ways by different geologists. Commonly, the chert appears to be composed entirely of crypto-crystalline or amorphous silica. To the north of the Buster Creek saddle the "chert" is a intermittently siliceous siltstone. My observations are mostly consistent with a marine to shallow marine origin for the chert. At the head of the South Fork of Bobtail Creek, the "chert" shows granular textures and some relic casts of pebbles and clasts



SCHEMATIC CROSS SECTION OF MARSHALL PLACER DISTRICT



SCHEMATIC CROSS SECTION OF KAKO PLACER DISTRICT




FIGURE 5

SCHEMATIC CROSS SECTION OF STUYAHOK PLACER DISTRICT

See geologic map of Lower Yukon gold belt for section location and explanation of rock types.

suggesting a partially recrystallized quartzite. However, this texture could also result from the re-silicification of a brecciated fine grained marine chert.

Fox, 1940, suggested that a large volume of this 'chert' was a silicified limestone. Some chert outcrops do weather in a distinctly fluted manner similar to carbonates. Fox also suggests that the conglomerate and breccia textures textures observed were a result of complete silicification of fine grained arenaceous sediments and recrystallized quartzite. RAA mapped these same rocks as a "siliceous tuff".

Rhyolites

The andesites and cherts have been intruded by a body of late Cretaceous to early Tertiary porphyritic albite rhyolite. The rhyolite is characterized by quartz eyes, clay-altered feldspar phenocrysts, and siliceous bands. Breccia, hypabyssal, and possibly siliceous dome, and proximal tuff textures are common in the rubblecrop. In addition to the areas exposed above the placer deposit, float and tailings found in the valley suggest that a large body of silica-rich rhyolite occurs immediately below the placer gravels.

The rhyolite that forms rubblecrop on the 901' ridge is sparsely porphyritic with 5% clear quartz eyes, rare to absent feldspar phenocrysts and minor relict sulfide grains, chiefly oxidized pyrite, in a white to buff aphanitic groundmass. Pebble clasts, and angular chert fragments are seen in some of the rhyolite. Alteration is light to moderate on the ridge and the argillic alteration and intense limonite alteration seen in the Buster Creek Saddle is not evident. Mafic grains are sparse and are generally altered to oxides but angular casts remain and groundmass staining is uncommon. Fragments from rubblecrop at the east end of line 300North displays siliceous banding and platy fracture.

It should be noted that although I believe the rhyolite is a Tertiary intrusion other geologists (RAA) have felt that this could be a marine bimodal volcanic sequence that is capable of hosting either Kuroko base metal or exhalite type gold mineralization.

The rhyolite alteration assemblage is characterized by clay, sericite and minor silica and iron carbonate. Limonite after sulfides and possibly after carbonate is common. Locally the alteration and silicification are so intense that the original composition and texture can not be identified.

Structure

The area has been extensively faulted by northeast and east-west trending faults. Sheared and brecciated rocks in rubble, the lack of continuous beds, and air photo identification of numerous lineaments all suggest unresolved structural complexities. The east-west fault which localizes the mineralized breccia above Buster Creek seems to be fairly high angle. Several areas appear to be intensely fractured on air photos. These areas surround the dome-like high points which occur along the ridge, Figure 4 and map labeled Geology of Kako Prospect.

Kako Placer

Buster Creek Gravels

The composition of the gravels is completely consistent with local derivation and short distances of transport. The Buster Creek placer was developed in the upper portion of an alluvial fan. Evidence of water table reduction/oxidation fronts is quite common but no relationship between stained gravels and the quality of pay has been established. The pay gravels are mostly rhyolite with minor chert and andesite. Our best estimate based on the extent of tailings is that 5 to 7 thousand ounces were won from Buster Creek.

Heavy Minerals

As can be expected from the surrounding country rocks there is only a small volume of "black sands" concentrated from the Kako placer. Placer concentrates from Buster Creek consist primarily of fine sized gold, cinnabar, garnet, and arsenopyrite. Minor amounts of scheelite, cassiterite, stibnite, ilmenite and magnetite are also reported. The minor volume of "black sands" and the composition of the concentrates is consistent with derivation from an epithermal gold system. Black sand concentrations increase in gravels derived from the Gemuk Group andesites.

Penz, who now uses a large jig system, confirms that a significant amount of the gold he recovers comes from nuggets on the one hand and minus 20 mesh on the other hand. This broad size distribution is typical for placers which are located close to their source.

The gold won from the Kako Placer is rough and coarse and rarely looks well traveled. Fine lace like structures thought to be box works or crystal casts are frequently observed on the nuggets. A small portion of the placer gold is found still attached to gangue minerals, typically quartz and sericite altered rhyolite. Rare pieces of gold attached to arsenopyrite have been reported. The gangue

minerals, angularity and shape of the gold, and the volume of cinnabar and stibnite in the placer concentrate are also consistent with a placer derived from an epithermal lode environment.

The fineness of the gold from the Kako placer is reported to be 818 (Smith, 1941). This low fineness is consistent with placer deposits near a lode source. Penz and past placer producers have reported the best yields from areas of rhyolite bedrock.

Bobtail Creek

The float in the upper section of the creek is composed chiefly of siliceous siltstone and chert. Rhyolite is a minor component of the float. No placer tailings were found above the 220' elevation where two tributary streams join Bobtail Creek from the north and south at the eastern edge of section 17. The valley floor is flat and willow choked at this elevation. Dragline overburden piles begin at the point where these tributaries enter the stream and continue downstream 2000 to 2500 feet where they encounter the upper end of a large beaver dam. Tailings between the dragline piles consist of 80-90% siltstone and shales, commonly iron-stained, with larger cobbles of chert and cherty tuffs. Crystal poor white rhyolite and andesitic volcanics are minor components of the tailings. No quartz was seen in the tailings. An estimated 2 to 4 thousand ounces was won from Bobtail Creek.

Placer Potential

Penz who is mining on Buster Creek has reserves on his claims sufficient for several years of operation. Additional placer reserves owned by Calista Corporation are known to exist on Buster and Bobtail Creek. Joesting, 1940, evaluated Buster Creek with a churn drill. He estimated that the average value for the remaining pay zones in Buster Creek was \$ 8.00 per yard (@ 400.00/oz) with \$4.00 cutoff grade. Joesting estimated that a minimum of 7 to 8 thousands ounces remained in Buster Creek, mostly on the lower portion of the creek (owned by Calista).

Significant geologic indicated placer reserves are thought to exist below and alongside the old tailings on Bobtail Creek and on adjacent Montezuma Creek. The upper portions of the Kuyukutuk River, north over the ridge from Buster and Bobtail Creeks, have also been reported to have favorable drill holes (Penz, 1989)

LODE TARGETS

The Kako Grid

In 1989, 265 pulps retained from RAA's 1974 soil grid program across Buster Saddle at Kako were re-analyzed by methods with greater

technical sophistication and lower detection levels than the original analyses done in 1974. The results defined two anomalous, open-ended zones within the grid area.

The soil sampling program conducted in 1989 by Calista was designed to expand the anomalous areas defined by RAA. A 6500 foot baseline oriented 45 degrees northeast was sampled at 50 foot intervals along Base Hill ridge west of the Kako placer mine. This line crossed contacts between all the lithologies present on the ridge. Two perpendicular lines trending northwest/southeast were also sampled to trace the extent of the anomalous rhyolites on either side of the Buster Saddle. The Calista soil sampling expanded the two anomalous zones discovered by RAA.

Table 1, found at the end of this report is a list of anomalous samples from the Kako Prospect. Figure 4 and the geochemical overlays accompanying the map Geology of the Kako Prospect show the locations of anomalous samples.

Rhyolite-Chert Contacts

The first anomalous area is a zone at the northern end of the grid near hill 901 with values ranging from 0.1 to 0.3 ppm gold and 2 to 3 ppm silver. Several anomalous samples cluster along rhyolite and chert or andesite contacts particularly where these contacts appear to intersect fault strikes.

In 1989 a 500' by 800' soil grid was sampled over this anomalous area. A horizon in the rocks mapped variously as chert, quartzite and siliceous tuff carries anomalous gold ranging from 110 to 260 ppb gold. A stream sediment sample taken from a small stream draining this chert yielded 360 ppb gold, Figure 4.

Half of the anomalous samples were in areas of rhyolite float, the other half were taken in soils with chert or cherty tuff float. A distinctive outcrop of fractured, bleached and highly limonite-stained chert is located at the top of hill 901. All samples from this area averaged 0.1 ppm gold. This outcrop is probably a cap or roof pendant on the rhyolite that is exposed as rubble north, south, and east of the chert. One thin section from a sample nearby the gold anomalies was determined to be a recrystallized sedimentary chert.

Buster Creek Breccia

The second zone is an area of limonite - quartz - rhyolite breccia, which is exposed in the saddle at the head of the South Fork of Buster Creek. The rock is vuggy, silicified, and has pervasive limonite staining. The rhyolite is pervasively altered by sericite and

clay.

This zone, which appears to be coincident and parallel with the trace of Buster Creek, carries gold values to 0.9 ppm, silver values up to 9.6 ppm, and arsenic levels greater than 2000 ppm. 32 of the re-analyzed samples were found to have gold values of .1 ppm or greater. This zone of brecciation is 30 feet wide and is traceable for 300 feet. The zone is lost in the brush below the saddle.

Because of the poor nature of the outcrop, it is difficult to determine grades across any width. However, soil and rock chip samples averaged together suggest that this zone **averages** 300 ppb gold, a little less than 2 ppm silver, and 500 ppm arsenic, see Table 1. Trenching in this area and further exploration along strike, particularly downstream toward the placer deposits is planned for the future.

A strong stream sediment anomaly has been identified below this occurrence. These samples which were taken above the placer tailings in the three streams which drain the east side of the Kako Grid area run 480, 200, 360, and 520 ppb gold, Figure 4 and Figure 5. This is one of the strongest stream sediment anomalies we have recorded in southwestern Alaska. Given the extent of the anomalies, the possibility of discovering higher grade mineralization in this area seems good.

Other Occurrences

In addition to the rhyolite breccia and the auriferous (?) chert reports of mineralized rhyolite dikes within the placer cuts and the occurrence of placer gold to the north in Bobtail Creek and to the west in the Kuyukutuk River suggest that other areas should be investigated.

Trace Element Geochemistry

Silver values are elevated on the steep southern slope of the 901' ridge. Values average between 1 and 2 ppm silver in a 400' by 800' area with rhyolite rubblecrop. Gold values are not correlative and are not anomalous in this area, however, arsenic and silver values show close correspondence. Silver anomalies are also present on the 901' soil grid. One highly anomalous sample had .542 ppm gold, 17.9 ppm silver, and 2000 ppm arsenic. Silver values in the 1-2 ppm range extend to line 700North.

Nearly the entire drainage is anomalous in arsenic. Arsenic is anomalous over the entire Buster Saddle and increases wherever rhyolite is found. The average arsenic level is about 200 ppm.

Arsenic values in brecciated or altered rhyolite commonly range from 500 to more than 2000 ppm. These levels are also present on the hill 901' chert with values of 667, 1384, and 2000 ppm over a one 100 foot interval. Gold and silver values often increase as arsenic increases. Sharp isolated jumps in arsenic content in areas of deep soil cover probably reflect small dikes or veins.

Lead, mercury, and antimony correlate with the occurrence of rhyolite porphyry. Mercury and antimony do not correlate with the presence or absence of gold. Despite the presence of cinnabar in the placer concentrates at Kako, the hardrock samples have weak values of mercury and antimony compared to the samples from the Stuyahok rhyolite.

Stuyahok History

Gold was discovered at Stuyahok in 1918 (Cobb, 1973). Little of the history of the operation is known. The mine ceased operation in the 1940's and the equipment was moved to Ruby. Total production in the area probably exceeds 20,000 ounces (Cobb, 1973).

In 1974 and 1975 geologists working for Resource Associates of Alaska (RAA) found anomalous gold, arsenic and mercury. Like Kako, many of the soil and stream sediments samples taken by RAA focused on base metal mineralization. Often no assays for gold or low temperature metals were run. RAA's assessment of the area was similar to the appraisal of the Kako property.

Calista selected the Stuyahok area in 1977 and leased the placer mine to the Chase Brothers in 1986. The five brothers and their father, who worked on the property in the early 1940's, have produced about 100 ounces. Calista terminated the lease in 1989 and the property is currently available for lease.

Calista accomplished some small scale mapping and sampling in 1985 and 1988. In 1989 careful sampling of the placer tailings resulted in the discovery of an area of gold bearing bedrock.

Stuyahok Geology

Lithologic Units

The ridge above the placer is underlain by coarse, angular andesite marine tuffs, flows, agglomerate and breccia. Felsic and siliceous tuff is occasionally interbedded with the andesite. On the north and east portions of the ridge, talus and rubble indicates that a quartz-feldspar porphyritic rhyolite with a very siliceous matrix intrudes the andesites, see Figure 5 and attached map Stuyahok Prospect Geology/Geochemistry.

